arXiv:1511.08671v1 [math.RA] 27 Nov 2015

A Note on Semigroup Algebras of Permutable Semigroups[1]

A. Nagy and M. Zubor

Abstract

Let S be a semigroup and $\mathbb{F}$ be a field. For an ideal J of the semigroup algebra $\mathbb{F}[S]$ of S over $\mathbb{F}$, let ϱ_J denote the restriction (to S) of the congruence on $\mathbb{F}[S]$ defined by the ideal J. A semigroup S is called a permutable semigroup if $\alpha \circ \beta = \beta \circ \alpha$ is satisfied for all congruences α and β of S. In this paper we show that if S is a semilattice or a rectangular band then $\varphi_{\{S;\mathbb{F}\}} : J \mapsto \varrho_J$ is a homomorphism of the semigroup $(Con(\mathbb{F}[S]); \circ)$ into the relations semigroup $(\mathcal{B}_S; \circ)$ if and only if S is a permutable semigroup.

For a non-empty set X, let $\mathcal{B}_X$ denote the set of all binary relation on a set X. It is known that $\mathcal{B}_X$ form a semigroup under the operation $\circ$ defined by the followong way: for arbitrary $\alpha, \beta \in \mathcal{B}_X$ and $x, y \in X$, $(x, y) \in \alpha \circ \beta$ if and only if there is an element $z \in X$ such that $(x, z) \in \alpha$ and $(z, y) \in \beta$. An algebraic structure A will said to be permutable if the congruences of A commute with each other (under the operation $\circ$). For arbitrary equivalences α and β of a set X, $\alpha \circ \beta$ is an equivalence on X if and only if $\alpha \circ \beta = \beta \circ \alpha$. Thus an algebraic stucture A is permutable if and only if the set $Con(A)$ of all congruences on A form a subsemigroup in the relation semigroup $(\mathcal{B}_A; \circ)$.

Let S be a semigroup and $\mathbb{F}$ be a field. For an ideal J of the semigroup algebra $\mathbb{F}[S]$, let ϱ_J denote the congruence on S which is the restriction of the congruence on $\mathbb{F}[S]$ defined by J. For an arbitrary congruence α on S, let $\mathbb{F}[\alpha]$ denote the kernel of the extended canonical homomorphishm $\mathbb{F}[S] \to \mathbb{F}[S/\alpha]$. By Lemma 5 of Chapter 4 of [5], for every semigroup S and every field $\mathbb{F}$, the mapping $\varphi_{\{S;\mathbb{F}\}} : J \mapsto \varrho_J$ is a surjective homomorphism of the semilattice $(Con(\mathbb{F}[S]); \wedge)$ onto the semilattice $(Con(S); \wedge)$ such that $\varrho_{\mathbb{F}[\alpha]} = \alpha$ for every congruence α on S. As a homomorphic image of a semigroup is also a semigroup, and $\alpha \circ \beta = \alpha \vee \beta$ is satisfied for every congruences α and β of a permutable semigroup, the assertions of the following lemma are obvious.

Lemma 1 *If S is a semigroup such that, for a field $\mathbb{F}$, $\varphi_{\{S;\mathbb{F}\}} : J \to \varrho_J$ is a homomorphism of the semigroup $(Con(\mathbb{F}[S]); \circ)$ into the semigroup $(\mathcal{B}_S; \circ)$ then S is a permutable semigroup. Moreover, if S is a permutable semigroup then $\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of $(Con(\mathbb{F}[S]); \circ)$ onto the semigroup $(Con(S); \circ)$ if and only if $\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of the semilattice $(Con(\mathbb{F}[S]); \vee)$ onto the lattice $(Con(S); \vee)$, that is, $ker_{\varphi_{\{S;\mathbb{F}\}}}$ is $\vee$-compatible.*

The next example shows that the converse of the first assertion of Lemma 1 is not true, in general; for a permutable semigroup S, the condition "$\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of $(Con(\mathbb{F}[S]); \circ)$ onto the semigroup $(Con(S); \circ)$" depends on the field $\mathbb{F}$.

[1]Keywords: permutable semigroup, semigroup algebra, lattice of congruences. Mathematics Subject Classification: 20M10, 20M25. National Research, Development and Innovation Office NKFIH, 115288

Example 1 *Let C_4, $\mathbb{F}_3$ and $\mathbb{F}_2$ denote the cyclic group of order 4, the 3-element field, and the 2-element field, respectively. It is known that every group is a permutable semigroup. Denote the elements of C_4 by $1, a, a^2, a^3$ (1 is the identity element). It is easy to see that $I = \mathrm{Span}(1+a^2, a+a^3)$ and $J = \mathrm{Span}(1+a, a+a^2, a^2+a^3)$ are ideals of $\mathbb{F}_3[C_4]$, and*

$$\varphi_{\{C_4;\mathbb{F}_3\}}(I) \vee \varphi_{\{C_4;\mathbb{F}_3\}}(J) =$$

$$\varrho_I \vee \varrho_J = \iota \vee \{(1, a^2), (a, a^3)\} \neq \omega_{C_4} = \varrho_{(I+J)} =$$

$$= \varrho_{(I \vee J)} = \varphi_{\{C_4;\mathbb{F}_3\}}(I \vee J).$$

Thus $ker_{\varphi_{\{C_4;\mathbb{F}_3\}}}$ is not $\vee$-compatible and so $\varphi_{\{C_4;\mathbb{F}_3\}}$ is not a homomorphism of $Con(\mathbb{F}_3[C_4]); \circ)$ onto the semigroup $(Con(C_4); \circ)$.

By computing the pricipal ideals of $\mathbb{F}_2[C_4]$ we get that the principal ideals of $\mathbb{F}_2[C_4]$ form a chain. As the dimension of $\mathbb{F}_2[C_4]$ is finite, from this it follows that $\mathbb{F}_2[C_4]$ has no other ideals. So $Con(\mathbb{F}_2[C_4])$ is the next (here α_{C_2} denotes the congruence of C_4 defined by C_2):



It is easy to see that $ker_{\varphi_{\{C_4;\mathbb{F}_2\}}}$ is $\vee$-compatible and so $\varphi_{\{C_4;\mathbb{F}_2\}}$ is a homomorphism of $Con(\mathbb{F}_2[C_4]); \circ)$ onto the semigroup $(Con(C_4); \circ)$.

By Lemma 1 and the above Example, it is a natural idea to find all couples $(S, \mathbb{F})$ of permutable semigroups S and fields $\mathbb{F}$, for which the mapping $\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of the semigroup $(\mathrm{Con}(\mathbb{F}[S]), \circ)$ onto the semigroup $(Con(S); \circ)$. In the next we show that if S is an arbitrary permutable semilattice or an arbitrary permutable rectangular band then, for an arbitrary field $\mathbb{F}$, $\varphi_{\{S;\mathbb{F}\}}$ satisfies the previous condition.

For notations and notions not defined here, we refer to [1, 4, 5, 6]

Theorem 1 *Let S be a permutable semilattice. Then, for an arbitrary field $\mathbb{F}$, $\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of the semigroup $(\mathrm{Con}(\mathbb{F}[S]), \circ)$ onto the semigroup $(Con(S); \circ)$.*

Proof. Assume that S is a permutable semilattice. Then, by Lemma 2 of [3], $|S| \leq 2$. We can consider the case when $|S| = 2$. Let

$$S = \{e, f\} \quad (e \neq f).$$

Then

$$e^2 = e, \quad \text{and} \quad f^2 = f.$$

We can suppose that

$$ef = fe = e.$$

It is clear that S has two congruences: ι_S and ω_S.

Let $\mathbb{F}$ be an arbitrary field. It is easy to see that

$$J_e = \{\alpha e : \ \alpha \in \mathbb{F}\} \quad \text{and} \quad J_{e-f} = \{\alpha(e-f) : \ \alpha \in \mathbb{F}\}$$

are proper ideals of $\mathbb{F}[S]$. As

$$dim(J_e) = dim(J_{e-f}) = 1,$$

the ideals J_e and J_{e-f} are minimal ideals of $\mathbb{F}[S]$. We show that the ideals of $\mathbb{F}[S]$ are

$$\{0\}, J_e, J_{e-f} \quad \text{and} \quad \mathbb{F}[S].$$

Let $J \neq \{0\}$ be a proper ideal of $\mathbb{F}[S]$. Clearly $dim(J) = 1$. Let

$$A = \alpha e + \beta f \in J$$

be a non-zero element. Then

$$(\alpha + \beta)e = e(\alpha e + \beta f) = eA \in J.$$

If $\alpha + \beta \neq 0$ then $J = J_e$. If $\alpha + \beta = 0$ then $J = J_{e-f}$.

Thus the ideals of $\mathbb{F}[S]$ are $\{0\}$, J_e, J_{e-f}, $\mathbb{F}[S]$. So $Con(\mathbb{F}[S])$ is the next:



It is a matter of checking to see that the $ker_{\varphi_{\{S;\mathbb{F}\}}}$-classes of $Con(\mathbb{F}[S])$ are $\{\{0\}, J_e\}$ and $\{J_{e-f}, \mathbb{F}[S]\}$. It is easy to see that $ker_{\varphi_{\{S;\mathbb{F}\}}}$ is $\vee$-compatible and so, by Lemma 1, $\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of the semigroup $(\mathrm{Con}(\mathbb{F}[S]), \circ)$ onto the semigroup $(Con(S); \circ)$. □

Corollary 1 *Let S be a semilattice. Then, for a field $\mathbb{F}$, $\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of the semigroup $(\mathrm{Con}(\mathbb{F}[S]), \circ)$ into the relation semigroup $(\mathcal{B}_S; \circ)$ if and only if $|S| \leq 2$.*

Proof. By Lemma 1, Theorem 1 and Lemma 2 of [3], it is obvious. □

Theorem 2 *Let $S = L \times R$ be a permutable rectangular band (L is a left zero semigroup, R is a right zero semigroup). Then, for an arbitrary field $\mathbb{F}$, $\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of the semigroup* $(\mathrm{Con}(\mathbb{F}[S]), \circ)$ *onto the semigroup* $(Con(S); \circ)$.

Proof. Let $\mathbb{F}$ be an arbitrary field and $S = L \times R$ be a permutable rectangular band. As a rectangular band satisfies the identity $axyb = ayxb$, that is, every rectangular band is a medial semigroup, Corollary 1.2 of [2] implies $|L| \leq 2$ and $|R| \leq 2$.

First consider the case when $|L| = 1$. Then S is isomorphic to the right zero semigroup R, and $|S| \leq 2$. We can suppose that $|S| = 2$. Let $S = \{e, f\}$ $(e \neq f)$. The congruences of S are ι_S and ω_S. We show that the ideals of $\mathbb{F}[S]$ are

$$\{0\}, J_{e-f} = \mathbb{F}[\omega_S] = \{\alpha(e-f) : \ \alpha \in \mathbb{F}\} \quad \text{and} \quad \mathbb{F}[S].$$

Let $J \neq \{0\}$ be an arbitrary ideal. Assume that there is an element

$$0 \neq \alpha_0 e + \beta_0 f \in J$$

for which

$$\alpha_0 + \beta_0 \neq 0$$

is satisfied. Then

$$(\alpha_0 + \beta_0)e = (\alpha_0 e + \beta_0 f)e \in J$$

and so

$$e = \frac{1}{\alpha_0 + \beta_0}(\alpha_0 + \beta_0)e \in J$$

from which we get $f = ef \in J$. Consequently

$$J = \mathbb{F}[S].$$

Next, consider the case when

$$\alpha_0 + \beta_0 = 0$$

is satisfied for every

$$A = \alpha_0 e + \beta_0 f \in J.$$

Then $\beta = -\alpha_0$ and so

$$A = \alpha_0 e + \beta_0 f = \alpha_0 e - \alpha_0 f = \alpha_0(e-f) \in J_{e-f}.$$

Consequently,

$$J \subseteq J_{e-f}.$$

As $dim(J_{e-f}) = 1$, the ideal J_{e-f} is minimal. Hence

$$J = J_{e-f}.$$

Thus the ideals of $\mathbb{F}[S]$ are $\{0\}$, J_{e-f} and $\mathbb{F}[S]$, indeed. So $Con(\mathbb{F}[S])$ is

$$\begin{array}{c} \mathbb{F}[S] \\ | \\ J_{e-f} \\ | \\ \{0\} \end{array}$$

It is a matter of checking to see that the $ker_{\varphi_{\{S;\mathbb{F}\}}}$-classes of $Con(\mathbb{F}[S])$ are $\{\{0\}\}$ and $\{J_{e-f}, \mathbb{F}[S]\}$. It is easy to see that $ker_{\varphi_{\{S;\mathbb{F}\}}}$ is $\vee$-compatible and so, by Lemma 1, $\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of the semigroup $(\mathrm{Con}(\mathbb{F}[S]), \circ)$ onto the semigroup $(Con(S); \circ)$.

If $|R| = 1$ then S is a left zero semigroup, and $|S| \leq 2$. We can prove, as in the previous part of the proof, that $ker_{\varphi_{\{S;\mathbb{F}\}}}$ is $\vee$-compatible.

Next, consider the case when $|L| = |R| = 2$. Let

$$L = \{a_1, a_2\}, \quad R = \{b_1, b_2\}.$$

Let α_L and α_R denote the kernels of the projection homomorphisms $S \mapsto L$ and $S \mapsto R$, respectively. The α_L-classes of S are

$$\{(a_1, b_1); (a_1, b_2)\} \quad \text{and} \quad \{(a_2, b_1); (a_2, b_2)\}.$$

The α_R-classes of S are

$$\{(a_1, b_1); (a_2, b_1)\} \quad \text{and} \quad \{(a_1, b_2); (a_2, b_2)\}.$$

It is easy to see that the congruences of S are ι_S, α_L, α_R and ω_S. We show that the ideals of $\mathbb{F}[S]$ are

$$\mathbb{F}[S], \mathbb{F}[\omega_S] = \{\sum_{i,j=1}^{2} \alpha_{i,j}(a_i, b_j) : \sum_{i,j=1}^{2} \alpha_{i,j} = 0\},$$

$$J_L = \mathbb{F}[\alpha_L], J_R = \mathbb{F}[\alpha_R], J_L \cap J_R, \{0\}.$$

We note that

$$dim(\mathbb{F}[\omega_S]) = 3, dim(J_L) = dim(J_R) = 2.$$

First we show that $J \subseteq \mathbb{F}[\omega_S]$ or $J = \mathbb{F}[S]$ for every ideal J of $\mathbb{F}[S]$. Let J be an arbitrary ideal of $\mathbb{F}[S]$. Assume that there is an element

$$A = \alpha_{1,1}(a_1, b_1) + \alpha_{1,2}(a_1, b_2) + \alpha_{2,1}(a_2, b_1) + \alpha_{2,2}(a_2, b_2) \in J$$

such that $A \notin \mathbb{F}[\omega_S]$, that is $\sum_{i,j=1}^{2} \alpha_{i,j} \neq 0$. Let $i, j \in \{1, 2\}$ be arbitrary elements. Then

$$(\sum_{i,j=1}^{2} \alpha_{i,j})(a_i, b_j) = (a_i, b_1)A(a_1, b_j) \in J.$$

As $\sum_{i,j=1}^{2} \alpha_{i,j} \neq 0$, we get $(a_i, b_j) \in J$ from which it follows that $S \subseteq J$. Consequently, $J = \mathbb{F}[S]$. Thus $\mathbb{F}[\omega_S]$ is the only maximal ideal of $\mathbb{F}[S]$.

Next we show that $J_L \cap J_R$ is the only ideal of $\mathbb{F}[S]$ whose dimension is 1. Let

$$A = \alpha_{1,1}(a_1, b_1) + \alpha_{1,2}(a_1, b_2) + \alpha_{2,1}(a_2, b_1) + \alpha_{2,2}(a_2, b_2) \in J_L \cap J_R$$

be an arbitrary element. As

$$(a_1, b_1)\ \alpha_L\ (a_1, b_2) \quad \text{and} \quad (a_2, b_1)\ \alpha_L\ (a_2, b_2),$$

we have

$$\alpha_{1,2} = -\alpha_{1,1} \quad \text{and} \quad \alpha_{2,2} = -\alpha_{2,1}.$$

As

$$(a_1, b_1)\ \alpha_R\ (a_2, b_1) \quad \text{and} \quad (a_1, b_2)\ \alpha_R\ (a_2, b_2),$$

we have

$$\alpha_{2,1} = -\alpha_{1,1} \quad \text{and} \quad \alpha_{2,2} = -\alpha_{1,2}.$$

Thus

$$A = \alpha_{1,1}((a_1, b_1) - (a_1, b_2) - (a_2, b_1) + (a_2, b_2))$$

for some $\alpha \in \mathbb{F}$. Consequently, the ideal $J_L \cap J_R$ is generated by

$$(a_1, b_1) - (a_1, b_2) - (a_2, b_1) + (a_2, b_2).$$

Hence the dimension of $J_L \cap J_R$ is 1.

To show that $J_L \cap J_R$ is the only ideal of $\mathbb{F}[S]$ whose dimension is 1, consider an ideal J of $\mathbb{F}[S]$ generated by an element

$$0 \neq B = \alpha_{1,1}(a_1, b_1) + \alpha_{1,2}(a_1, b_2) + \alpha_{2,1}(a_2, b_1) + \alpha_{2,2}(a_2, b_2).$$

Then $J \subset \mathbb{F}[\omega_S]$ and

$$(a_1, b_1)B = (\alpha_{1,1} + \alpha_{2,1})(a_1, b_1) + (\alpha_{1,2} + \alpha_{2,2})(a_1, b_2) \in J.$$

Thus there is a coefficient $\xi \in \mathbb{F}$ such that

$$(a_1, b_1)B = \xi B.$$

Assume $\xi \neq 0$. Then $\alpha_{2,1} = \alpha_{2,2} = 0$ and so

$$B = \alpha_{1,1}(a_1, b_1) + \alpha_{1,2}(a_1, b_2).$$

From

$$(a_2, b_2)B = \alpha_{1,1}(a_2, b_1) + \alpha_{1,2}(a_2, b_2) \in J$$

we can conclude that $\alpha_{1,1} = \alpha_{1,2} = 0$ and so $B = 0$. This is a contradiction. Hence $\xi = 0$. Thus

$$B = \alpha_{1,1}(a_1, b_1) + \alpha_{1,2}(a_1, b_2)\alpha_{1,1}(a_2, b_1) - \alpha_{1,2}(a_2, b_2).$$

As

$$B(a_1, b_1) = (\alpha_{1,1} + \alpha_{1,2}((a_1, b_1) - (a_2, b_1)) \in J,$$

we get $B(a_1, b_1) = \tau B$ for some $\tau \in \mathbb{F}$. Assume $\tau \neq 0$. Then $\alpha_{1,2} = 0$ and so

$$B = \alpha_{1,1}(a_1, b_1) - \alpha_{1,1}(a_2, b_1).$$

From

$$B(a_2, b_2) = \alpha_{1,1}((a_1, b_2) - (a, 2, b_2)) \in J$$

we can conclude that $\alpha_{1,1} = 0$ and so $B = 0$. This is a contradiction. Hence $\tau = 0$. Thus $\alpha_{1,2} = -\alpha_{1,1}$ and so

$$B = \alpha_{1,1}((a_1, b_1) - (a_1, b_2) - (a_2, b_1) + (a_2, b_2)) \in J_L \cap J_R.$$

As $J \neq \{0\}$ and $J_L \cap J_R$ is a minimal ideal of $\mathbb{F}[S]$, we get

$$J = J_L \cap J_R,$$

that is, $J_L \cap J_R$ is the only ideal of $\mathbb{F}[S]$ whose dimension is 1.

As $dim(J_R + J_L) > dim J_R$ and $\mathbb{F}[\omega_S] \supseteq J_R + J_L$, we have

$$J_R + J_L = \mathbb{F}[\omega_S].$$

Let J be an arbitrary ideal of $\mathbb{F}[S]$ which differs from all of the ideals $\mathbb{F}[S], \mathbb{F}[\omega_S], J_L, J_R, J_L \cap J_R, \{0\}$. Then $J \subset \mathbb{F}[\omega_S]$ and $dim(J) = 2$.

If $J \cap J_L = \{0\}$ then $dim(J + J_L) = 4$ which contradicts $J + J_L \subseteq \mathbb{F}[\omega_S]$. Hence $dim(J \cap J_L) = 1$ and so $J_L \cap J_R = J \cap J_L$. From this we get $J \cap J_R = J_L \cap J_R$. Recall that $C = (a_1, b_1) - (a_1, b_2) - (a_2, b_1) + (a_2, b_2)$ generates the ideal $J_L \cap J_R$. Let A be an arbitrary element of $J \setminus (J_L \cap J_R)$. Then A and C are linearly independent. So

$$A = \alpha(a_1, b_1) + \beta(a_1, b_2) + \gamma(a_2, b_1) + (-\alpha - \beta - \gamma)(a_2, b_2).$$

If $\alpha = -\gamma$ then $A \in J_R$ which is a contradiction. Thus $\alpha \neq -\gamma$. Then

$$(a_1, b_1)A = (\alpha + \gamma)((a_1, b_1) - (a_1, b_2)) \in J_L.$$

As J is an ideal and $A \in J$ we have

$$(a_1, b_1)A \in J \cap J_L = J_L \cap J_R.$$

It means $\alpha = -\gamma$ which is also contradiction. Thus $Con(\mathbb{F}[S])$ is the next:



It is a matter of checking to see that the $ker_{\varphi_{\{S;\mathbb{F}\}}}$-classes of $Con(\mathbb{F}[S])$ are $\{\{0\}, J_L \cap J_R\}, \{J_L\}, \{J_R\}$ and $\{\mathbb{F}[\omega_S], \mathbb{F}[S]\}$. It is easy to see that $ker_{\varphi_{\{S;\mathbb{F}\}}}$ is $\vee$-compatible and so, by Lemma 1, $\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of the semigroup $(\mathrm{Con}(\mathbb{F}[S]), \circ)$ onto the semigroup $(Con(S); \circ)$. □

Corollary 2 *Let $S = L \times R$ be a rectangular band. Then, for a field $\mathbb{F}$, $\varphi_{\{S;\mathbb{F}\}}$ is a homomorphism of the semigroup $(\mathrm{Con}(\mathbb{F}[S]), \circ)$ into the relation semigroup $(\mathcal{B}_S; \circ)$ if and only if $|L|, |R| \leq 2$.*

Proof. By Lemma 1, Theorem 2 and Corollary 1.2 of [2], it is obvious. □

References

[1] Clifford, A.H. and G.B. Preston, *The Algebraic Theory of Semigroups*, Amer. Math. Soc., Providence, R.I., I(1961)

[2] Bonzini, C. and A. Cherubini, *Medial permutable semigroups*, Proc. Coll. Math. Soc. János Bolyai, 39. Semigroups, Szeged (Hungary), 1981, 21-39

[3] Hamilton, H., *Permutability of congruences on commutative semigroups*, Semigroup Forum, 10(1975), 55-66

[4] Nagy, A., *Special Classes of Semigroups*, Kluwer Academic Publishers, Dordrecht/Boston/London, 2001

[5] Okniński, J., *Semigroup Algebras*, Marcel Dekker, Inc., New York/Basel/Hong Kong, 1991

[6] Petrich, M., *Lectures in Semigroups*, Akademie-Verlag-Berlin, 1977

Department of Algebra
Institute of Mathematics
Budapest University of Technology and Economics
1521 Budapest, Pf. 91
e-mail (A. Nagy): nagyat@math.bme.hu
e-mail (M. Zubor): zuborm@math.bme.hu